

### **Technical Data Sheet**

# Reinforced Clostridial Agar (RCA)

Ordering number: 1.05410.0500

Medium proposed by BARNES and INGRAM (1956) for the cultivation and enumeration of clostridia, other anaerobes and facultative microorganisms in foodstuffs and other materials.

MUNOA and PARES (1988) developed a Bifidobacterium Iodoacetate Medium (BIM-25) on the basis of Clostridial Agar for the selective cultivation and differentiation of Bifidobacterium species.

#### **Mode of Action**

This culture medium is free from inhibitors and contains cysteine as a reducing agent. According to HIRSCH and GRINSTED (1954), Polymyxin B can be added to inhibit Gramnegative bacteria.

### Typical Composition (g/L)

Reinforced Clostridial Agar		
Meat extract*	10.0	
Peptone from casein	10.0	
Yeast extract	3.0	
D(+)glucose	5.0	
Starch	1.0	
Sodium chloride	5.0	
Sodium acetate	3.0	
L-cysteinium chloride	0.5	
Agar-agar**	12.5	

<sup>\*</sup> Meat extract is equivalent to the term beef extract.

MERCK

1.05410.0500

<sup>\*\*</sup>Agar-agar is equivalent to other different terms of agar.

#### **Preparation**

Suspend 50 g/litre, if desired dispense into test tubes, autoclave (15 min at 121 °C). If required, cool to 45-50 °C and add 0.02 g Polymyxin B/litre in form of a filter-sterilized aqueous solution.

pH:  $6.8 \pm 0.2$  at 25 °C.

The medium in the tubes or Petri dishes is clear and yellowish-brown.

### **Experimental Procedure and Evaluation**

Prepare stab cultures of the sample material in test tubes or use the pour-plate technique.

Incubation: 24-48 hours at an optimal temperature (e.g. 35 °C) under anaerobic conditions (e.g. Anaerocult® A, Anaerocult® A mini, or Anaerocult® P).

Count the colonies and, if necessary, perform additional tests.

#### **Storage**

Store at +15 °C to +25 °C, dry and tightly closed. Do not use clumped or discolored medium.

Protect from UV light (including sun light). For in vitro use only.

## **Quality Control**

Control strains	Growth after 48h at 35°C*	
Clostridium bifermentans ATCC® 19299	good / very good	
Clostridium novyi ATCC® 17861	good / very good	
Clostridium sporogenes ATCC® 19404 (WDCM 00008)	good / very good	
Clostridium perfringens ATCC® 13124 (WDCM 00007)	good / very good	
Clostridium perfringens ATCC® 10543	good / very good	
Escherichia coli ATCC® 25922 (WDCM 00013)	good / very good	
Bacillus cereus ATCC® 11778 (WDCM 00001)	good / very good	

<sup>\*</sup>QC strains were mixed in tubes with liquefied RCA, then solidified to generate anaerobic conditions and placed in the incubator

Please refer to the actual batch related Certificate of Analysis.



1.05410.0500 Page 2 of 3

#### Literature

BARNES, E.M., a. INGRAM, M.: The effect of redox potential on the grown Clostridium welchii strain isolated from horse muscle. – **J. Appl. Bact., 19**; 177-178 (1956).

HIRSCH, A., a. GRINSTED, E.: Methods for the growth and enumeration of anaerobic sporeformers from cheese, with observations on the effect of nisin. – **J. Dairy Res., 21**;101-110 (1954).

MUNOA, F.J., a. PARES, R.: Selective medium for isolation and enumeration of Bifidobacterium spp. – **Appl. Environ. Microbiol., 54**; 1715-1718 (1988).

#### **Ordering Information**

Product	Cat. No.	Pack size
Reinforced Clostridial Agar (RCA)	1.05410.0500	500 g
Anaerobic jar	1.16387.0001	1 ea
Anaeroclip®	1.14226.0001	1 x 25
Anaerocult® A	1.13829.0001	1 x 10
Anaerocult® A mini	1.01611.0001	1 x 25
Anaerocult® P	1.13807.0001	1 x 25
Anaerotest®	1.15112.0001	1 x 50
Plate basket	1.07040.0001	1 ea
Polymyxin-B-sulfate	Calbiochem 5291-1GM	1 g

Merck KGaA

Frankfurter Strasse 250 64293 Darmstadt, Germany Fax: +49 (0) 61 51 / 72-60 80 Find contact information for your country at: www.merckmillipore.com/offices

For Technical Service, please visit: www.merckmillipore.com/techservice

For more information, visit

www.merckmillipore.com/biomonitoring

Merck, Millipore, and Sigma-Aldrich are trademarks of Merck KGaA, Darmstadt, Germany or its affiliates. Detailed information on trademarks is available via publicly accessible resources. © 2019 Merck KGaA, Darmstadt, Germany and/or its affiliates. All Rights Reserved.



The life science business of Merck operates as

